

[54] CERTAIN UNSYMMETRICAL QUINOLINYL ETHERS HAVING ANTI-INFLAMMATORY AND ANTI-ALLERGIC ACTIVITY

[75] Inventors: John H. Musser, Malvern, Pa.; Utpal R. Chakraborty, Orangeburg, N.Y.

[73] Assignee: USV Pharmaceutical Corporation, Tuckahoe, N.Y.

[*] Notice: The portion of the term of this patent subsequent to Jan. 28, 2003 has been disclaimed.

[21] Appl. No.: 810,868

[22] Filed: Dec. 19, 1985

Related U.S. Application Data

[63] Continuation-in-part of Ser. No. 666,430, Oct. 30, 1984, abandoned, which is a continuation-in-part of Ser. No. 530,811, Sep. 9, 1983, Pat. No. 4,567,184, which is a continuation-in-part of Ser. No. 445,876, Dec. 1, 1982, abandoned.

[51] Int. Cl.⁴ C07D 215/22; C07D 217/24

[52] U.S. Cl. 546/152; 546/122; 546/149; 546/153; 546/176; 546/180; 546/339; 546/344; 544/250; 544/283; 544/345; 544/353; 544/385; 544/408; 549/365

[58] Field of Search 546/152, 149, 153, 180

[56] References Cited

U.S. PATENT DOCUMENTS

2,862,956 12/1958 Gundel et al. 560/73
4,360,700 11/1982 Melvin, Jr. 568/644

FOREIGN PATENT DOCUMENTS

0113587 7/1984 European Pat. Off. 546/344
2253511 7/1975 France 568/336
175293 12/1976 New Zealand 568/636
184606 6/1979 New Zealand 558/414
200031 11/1984 New Zealand 546/153

OTHER PUBLICATIONS

Boots, et al., Journal of Pharmaceutical Sciences, vol. 62, No. 6, Jun. 1973.

European Search Report, EP 83 11 2031.

Chemical Abstracts, vol. 94, 1981, p. 709, 174893.

Primary Examiner—Alan L. Rotman

Attorney, Agent, or Firm—Scully, Scott, Murphy & Presser

[57] ABSTRACT


Compounds of the formula:





and salts thereof, wherein

Ar₁ is a nitrogen, sulfur or oxygen heterocyclic ring;
Ar is a phenyl ring or a nitrogen, oxygen or sulfur heterocyclic ring;Ar and Ar₁ may be fully substituted or less than fully substituted with H, CH₃, lower alkyl, aryl, aralkyl, halo, hydroxy, lower alkoxy, CF₃, carboxy, alkylcarboxy, arylcarboxy, alkylcarbalkoxy, alkanoyl, formyl, oxo, nitrilo, amino, aminoalkyl, alkylamine, carboxamide, aryloxy, nitro, sulfonyl, sulfonamide, thio, alkylthio, hydroxyalkyl or oxyalkylcarbalkoxy;

X=



of up to 2 carbon atoms in the principal chain and up to a total of 4 carbon atoms,

Z is an alkylene chain containing up to 10 carbon atoms in the principal chain and a total of up to 12 carbon atoms and from 0 to 2 double bonds and the said alkylene chain may be attached to Ar through an oxygen, sulfur or amino nitrogen atom, and when n'=2, one of the R substituents may be halo; when n'=1, R is a substituent attached to one of the carbon atoms of Z selected from the group consisting of =O, OR₃, SR₃, N(R₂)₂ and R₁, —COR₄ and when n'=2 one R is previously defined and the additional R is a substituent attached to one of the carbon atoms of Z selected from the group consisting of =O, OR₃, SR₃, N(R₂)₂, —COR₄, lactone and halo;R₁ is H or CH₃;R₂ is H, lower alkyl, aryl or aralkyl;R₃ is H, lower alkyl, lower alkanoyl, aryl, aralkyl or substituted aryl in which the substituent is halo, lower alkyl or lower alkoxy;R₄ is OR₂ or N(R₂)₂;

n=0 or 1;

n'=1 to 7; and

n''=0, 1 or 2.

36 Claims, No Drawings